

Here to help, from start to finish

Shawn

8 YEARS OLD
on VOXZOGO since
4 years old as part
of a clinical trial



"I'm around for product education and support, as long as a family needs while their child is on VOXZOGO."

— Sonya

VOXZOGO is [the only/a] treatment FDA approved from birth to increase linear growth for children with achondroplasia and open growth plates.

IMPORTANT SAFETY INFORMATION

VOXZOGO may cause serious side effects including a temporary decrease in blood pressure in some patients. To reduce the risk of a decrease in blood pressure and associated symptoms (dizziness, feeling tired, or nausea), patients should eat a meal and drink 8 to 10 ounces of fluid within 1 hour before receiving VOXZOGO.

Please see additional safety information throughout as well as the included full **Prescribing Information** and **Patient Information**.



One-to-one support for you and your family:

- **Tailoring product and injection education** to your needs—available to extended family and other caregivers as well
- **Helping you to find an injection routine** that works for your family through the entire duration of your child's time on treatment
- **Supporting you through the specialty pharmacy process** and reminders for product refills
- **Offering flexible options** to meet you in person and/or virtually, by email, phone, or text



Case and Field Reimbursement Managers help navigate coverage by:

- **Helping with the insurance process** and sharing coverage options
- **Identifying financial programs**, such as co-pay assistance, if you're an eligible commercially insured patient*
- **Offering guidance** on insurance changes
- **Ensuring the specialty pharmacy** receives your prescription
- **Educating your doctor on coverage** requirements for continued access to therapy

Clinical Coordinator: _____

Phone: _____ **Email:** _____

*Terms and Conditions apply. Valid only for patients with commercial prescription insurance coverage who have a valid prescription for an FDA-approved indication and who meet additional eligibility criteria. Not valid for prescriptions reimbursed, in whole or in part, by any federal, state, or government-funded insurance programs (for example, Medicare, Medicare Advantage, Medigap, Medicaid, VA, DoD, or TRICARE) or where prohibited by law or by the patient's health insurance provider. If at any time a patient begins receiving prescription drug coverage under any federal, state, or government-funded healthcare program, the patient will no longer be able to use the program and patient must notify BioMarin RareConnections at 1-866-906-6100 to stop participation. Patients residing in or receiving treatment in certain states may not be eligible for some or all of the program elements. Patients may not seek reimbursement for value received from the program from any third-party payers. Additional restrictions may apply. Offer subject to change or discontinuance without notice. This assistance offer is not health insurance. See BioMarin-copay-terms.com for full Terms and Conditions.

IMPORTANT SAFETY INFORMATION

What is VOXZOGO used for?

- VOXZOGO is a prescription medicine used to increase linear growth in children with achondroplasia and open growth plates (epiphyses).
- VOXZOGO is approved under accelerated approval based on an improvement in annualized growth velocity. Continued approval may be contingent upon verification and description of clinical benefit in confirmatory trials.

What is the most important safety information about VOXZOGO?

- VOXZOGO may cause serious side effects including a temporary decrease in blood pressure in some patients. To reduce the risk of a decrease in blood pressure and associated symptoms (dizziness, feeling tired, or nausea), patients should eat a meal and drink 8 to 10 ounces of fluid within 1 hour before receiving VOXZOGO.

What are the most common side effects of VOXZOGO?

- The most common side effects of VOXZOGO include injection site reactions (including redness, itching, swelling, bruising, rash, hives, and injection site pain), high levels of blood alkaline phosphatase shown in blood tests, vomiting, joint pain, decreased blood pressure, and stomach ache. These are not all the possible side effects of VOXZOGO. Ask your healthcare provider for medical advice about side effects, and about any side effects that bother the patient or that do not go away.

How is VOXZOGO taken?

- VOXZOGO is taken daily as an injection given under the skin, administered by a caregiver after a healthcare provider determines the caregiver is able to administer VOXZOGO. Do not try to inject VOXZOGO until you have been shown the right way by your healthcare provider. VOXZOGO is supplied with Instructions for Use that describe the steps for preparing, injecting, and disposing VOXZOGO. Caregivers should review the Instructions for Use for guidance and any time they receive a refill of VOXZOGO in case any changes have been made.
- Inject VOXZOGO 1 time every day, at about the same time each day. If a dose of VOXZOGO is missed, it can be given within 12 hours from the missed dose. After 12 hours, skip the missed dose and administer the next daily dose as usual.
- The dose of VOXZOGO is based on body weight. Your healthcare provider will adjust the dose based on changes in weight following regular check-ups.
- Your healthcare provider will monitor the patient's growth and tell you when to stop taking VOXZOGO if they determine the patient is no longer able to grow. Stop administering VOXZOGO if instructed by your healthcare provider.

What should you tell the doctor before or during taking VOXZOGO?

- Tell your doctor about all of the patient's medical conditions including
 - If the patient has heart disease (cardiac or vascular disease), or if the patient is on blood pressure medicine (anti-hypertensive medicine).
 - If the patient has kidney problems or renal impairment.
 - If the patient is pregnant or plans to become pregnant. It is not known if VOXZOGO will harm the unborn baby.
 - If the patient is breastfeeding or plans to breastfeed. It is not known if VOXZOGO passes into breast milk.
- Tell your doctor about all of the medicines the patient takes, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

You may report side effects to BioMarin at **1-866-906-6100**. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call **1-800-FDA-1088**.

Please see additional safety information in the full Prescribing Information and Patient Information.